



Epidemiology and Response Division

NEW MEXICO INFLUENZA SURVEILLANCE UPDATE from the Epidemiology and Response Division of the New Mexico Department of Health (NMDOH) Weekly Report ending January 15, 2005

Effective immediately, the Public Health Order restricting the use of influenza vaccine has been rescinded.

Summary of Influenza Activity in New Mexico for Week Ending January 15, 2005:

- Eighteen of the 18 sentinel sites reported a total of 4371 patient visits, of which 35(0.80%) were for an influenza-like illness¹. The previous week ending January 8 reported 0.45 % influenza-like illness.
- NMDOH received reports of 30 patients with positive influenza (13 influenza A, 13 influenza B, 4 indistinguishable²) tests using rapid testing. There were two reports of positive influenza A culture (one of which was subtyped as H3), and 3 reports of influenza B (two of which have been subtyped as Sichuan-like).
- NMDOH reported the state influenza activity as “**SPORADIC**” to the Centers for Disease Control and Prevention (CDC) (see table below for definitions).

Laboratory Activity in NM:

- To date this season, there has been four influenza B virus isolates (3 subtyped as Sichuan-like) and four influenza A virus isolates* (three subtyped as H3) identified by culture at NMDOH Scientific Laboratory Division (SLD). After the first report of culture-confirmed influenza for the season, influenza activity reported to the CDC includes results from influenza rapid testing, fluorescent antibody (DFA) methods, or cultures.
- For the week ending January 15, 2005, fifteen clinical laboratories reported performing 260 rapid or DFA tests, of which 13(5.0%) were positive for influenza A, 13(5.0%) were positive for influenza B and 4(1.54%) were positive for indistinguishable² influenza type.
- Since October 24, 2004, NMDOH has received reports of 30(2.46%) positive influenza A tests, 32(2.62%) positive influenza B tests and 4(0.33%) indistinguishable² positive influenza out of 1221 rapid tests performed at 16 clinical laboratories.

*These cases may also be counted among the rapid test positive results.

¹ Influenza-like Activity (ILI) is defined as Fever ($\geq 100^{\circ}\text{F}$ [37.8°C], oral or equivalent) AND cough and/or sore throat in absence of a KNOWN cause other than influenza.

² Some rapid influenza tests cannot differentiate between types A and B.

Influenza-related Pediatric Mortality

As of the week ending January 8, 2005, one case of influenza-associated pediatric death have been reported to nationally to CDC. This report came from the Bureau of Health in Maine. No cases have been reported to the NMDOH.

Flu Activity in the Region

For the week ending January 8, 2005 (the most recent data available), influenza activity was reported as “widespread” by Texas, “regional” by Colorado, Idaho and Nevada, “local” by Utah and “sporadic” in Arizona, Montana, New Mexico and Wyoming. There were 10 reports of influenza A (H3N2) virus, 20 influenza A that were not subtyped and 17 influenza B viruses in the Mountain region (Arizona, Colorado, Idaho, Montana, Nevada, New Mexico, Utah and Wyoming).²

National Flu Surveillance and Laboratory Activity

For the week ending January 8, 2005, 510 (16.8 %) of 3,040 specimens tested for influenza viruses were positive. Of these one was an influenza A (H1N1), 69 were influenza A (H3N2) virus, 383 were influenza A that were not subtyped, and 57 were influenza B viruses. Nationwide 2.5% of patient visits to U.S. sentinel providers were due to influenza-like-illness. Eight states reported widespread activity, 10 states reported regional activity, 10 states reported local activity and 20 states reported sporadic activity nationally. More information on national surveillance can be found at <http://www.cdc.gov/flu/weekly/>.

This information is collected by the Infectious Disease Epidemiology Bureau, Epidemiology Response Division, NMDOH. For questions, please call 505-827-0006. For more information on influenza go to the NMDOH web page: <http://www.health.state.nm.us/flu/> or the CDC web page: <http://www.cdc.gov/ncidod/diseases/flu/fluvirus.htm>

Activity Level	ILI activity*/Outbreaks		Laboratory data
No activity	Low	And	No lab confirmed cases [†]
Sporadic	Not increased	And	Isolated lab-confirmed cases
	OR		
Local	Not increased	And	Lab confirmed outbreak in one institution [‡]
	Increased ILI in 1 region**; ILI activity in other regions is not increased	And	Recent (within the past 3 weeks) lab evidence of influenza in region with increased ILI
OR			

² All data are preliminary and change as more reports are received after the end of the reporting week.

	2 or more institutional outbreaks (ILI or lab confirmed) in 1 region; ILI activity in other regions is not increased	And	Recent (within the past 3 weeks) lab evidence of influenza in region with the outbreaks; virus activity is no greater than sporadic in other regions
Regional (doesn't apply to states with ≤ 4 regions)	Increased ILI in ≥ 2 but less than half of the regions	And	Recent (within the past 3 weeks) lab confirmed influenza in the affected regions
	OR		
	Institutional outbreaks (ILI or lab confirmed) in ≥ 2 and less than half of the regions	And	Recent (within the past 3 weeks) lab confirmed influenza in the affected regions
Widespread	Increased ILI and/or institutional outbreaks (ILI or lab confirmed) in at least half of the regions	And	Recent (within the past 3 weeks) lab confirmed influenza in the state.


* ILI activity can be assessed using a variety of data sources including sentinel providers, school/workplace absenteeism, and other syndromic surveillance systems that monitor influenza-like illness.

† Lab confirmed case = case confirmed by rapid diagnostic test, antigen detection, culture, or PCR. Care should be given when relying on results of point of care rapid diagnostic test kits during times when influenza is not circulating widely. The sensitivity and specificity of these tests vary and the predictive value positive may be low outside the time of peak influenza activity. Therefore, a state may wish to obtain laboratory confirmation of influenza by testing methods other than point of care rapid tests for reporting the first laboratory confirmed case of influenza of the season.

‡ Institution includes nursing home, hospital, prison, school, etc.

**Region: population under surveillance in a defined geographical subdivision of a state. A region could be comprised of 1 or more counties and would be based on each state's specific circumstances. Depending on the size of the state, the number of regions could range from 2 to approximately 12. The definition of regions would be left to the state but existing state health districts could be used in many states. Allowing states to define regions would avoid somewhat arbitrary county lines and allow states to make divisions that make sense based on geographic population clusters. Focusing on regions larger than counties would also improve the likelihood that data needed for estimating activity would be available.

Percentage of Visits for Influenza-like Illness
Reported by Sentinel Providers 2004 - 2005



Percentage of Visits for Influenza-like Illness
Reported by Sentinel Providers 2003 - 2004

