

# **New Mexico Pregnancy Risk Assessment (PRAMS)**

Multiyear Data Tables, Birth Years 1998-2003

NM PRAMS

nm.prams@state.nm.us

Telephone 505-476-8895

NM PRAMS, Family Health Bureau

2040 South Pacheco St.

Santa Fe, NM 87505

<http://www.health.state.nm.us/phd/prams/>

## **Suggested citation:**

NM Pregnancy Risk Assessment and Monitoring System (PRAMS).

Multiyear data tables, birth years 1998-2003. Santa Fe, NM: NM Department of Health, June 2005.

## **Funding for NM PRAMS is provided by:**

NM Department of Health

The Centers for Disease Control and Prevention (CDC) Cooperative Grant

U50/CCU613632

Federal Title V Maternal and Child Health Block Grant

Federal Family Preservation and Support Project of the NM Children, Youth and Families Department

Medical Assistance Division of the NM Department of Human Services.

## **New Mexico Pregnancy Risk Assessment (PRAMS) Multiyear Data Tables, Birth Years 1998-2003**

PRAMS surveys new mothers within two to six months after delivery, asking about experiences, attitudes, and behaviors related to infant and maternal health. Data are used to help plan and evaluate programs, inform policymakers, and educate healthcare providers and the public.

In these tables, questions follow same order as the table of contents (introductory section) of the NM PRAMS Surveillance Report, Birth Years 2001-2002, which is available on line or in print from NM PRAMS. The Surveillance Report includes the public health importance of each topic, actions addressing the issues, and the methodology of the PRAMS project.

Topics include preconception health (folic acid, intention of pregnancy, family planning); maternal behaviors and experiences, (alcohol and tobacco use, infant's smoke exposure, physical abuse by a partner, maternal weight problem and diabetes); care during the prenatal and postpartum periods (timely prenatal care, topics discussed during prenatal care, payer of care, health-related services); infant care (breastfeeding, infant sleep position, well-child care); and food sufficiency.

The lower and upper limits of each confidence interval (95% CI) appear in the columns labeled "lower" and "upper," respectively. If the figure in the column, " $\pm$ " is subtracted from or added to the "percent" figure, this gives the limits of the lower or upper 95% CI. When comparing estimates, if the 95% CI's do not overlap, we assume that there is a difference. However, if there is a small overlap, we need to do statistical tests to determine whether or not there is a difference.

Potential sources of bias include non-response, excluding non-resident and out-of-state births, and dropping Navajo women living on reservations from 2 months in 2002 and 5 months in 2003.

## NM Pregnancy Risk Assessment Monitoring System, birth years 1998–2003

These tables follow the order of the NM PRAMS Surveillance Report, birth years 2001-2002, Table of Contents. The Surveillance Report provides narrative with public health importance and actions addressing issues.

Overall number of respondents, n=9436. Item response available on request. Strikethrough indicates large confidence interval (>15 percentage points) or relative error (>0.3). Unless indicated, responses are generalized to all mothers with recent live birth.

	Year of birth	Percent	±
<b>Preconception issues</b>			
<b>% who were aware that folic acid can help prevent birth defects</b>			
	1998-2003	<b>64.3</b>	1.1
	1998	<b>66.3</b>	3.4
	1999	<b>68.9</b>	2.6
	2000	<b>63.1</b>	2.5
	2001	<b>65.5</b>	2.6
	2002	<b>63.5</b>	2.6
	2003	<b>58.4</b>	2.8
<b>% who used a multivitamin daily during the month before pregnancy</b>			
	2000-2003	<b>23.5</b>	1.1
	2000	<b>23.9</b>	2.2
	2001	<b>22.2</b>	2.2
	2002	<b>22.4</b>	2.2
	2003	<b>25.4</b>	2.5
<b>% who did not use a multivitamin at all during the month before pregnancy</b>			
	2000-2003	<b>61.7</b>	1.3
	2000	<b>63.0</b>	2.5
	2001	<b>62.1</b>	2.6
	2002	<b>61.0</b>	2.5
	2003	<b>60.8</b>	2.8
<b>% who intended the pregnancy</b>			
	1998-2003	<b>56.6</b>	1.2
	1998	<b>57.1</b>	3.7
	1999	<b>56.4</b>	2.9
	2000	<b>56.4</b>	2.6
	2001	<b>57.7</b>	2.7
	2002	<b>55.8</b>	2.6
	2003	<b>56.1</b>	2.8
<b>% who had mistimed pregnancy (wanted later)</b>			
	1998-2003	<b>32.7</b>	1.1
	1998	<b>30.9</b>	3.5
	1999	<b>32.4</b>	2.7
	2000	<b>33.1</b>	2.4
	2001	<b>31.6</b>	2.5
	2002	<b>33.8</b>	2.5
	2003	<b>34.0</b>	2.7
<b>% who had unwanted pregnancy (not wanted then or ever)</b>			
	1998-2003	<b>10.8</b>	0.7
	1998	<b>12.0</b>	2.5
	1999	<b>11.2</b>	1.8
	2000	<b>10.5</b>	1.6
	2001	<b>10.8</b>	1.7
	2002	<b>10.5</b>	1.6
	2003	<b>9.9</b>	1.8

	Year of birth	Percent	±
<b>% who had unintended pregnancy (mistimed + unwanted)</b>			
	1998-2003	<b>43.4</b>	1.2
	1998	<b>42.9</b>	3.7
	1999	<b>43.6</b>	2.9
	2000	<b>43.6</b>	2.6
	2001	<b>42.3</b>	2.7
	2002	<b>44.2</b>	2.6
	2003	<b>44.0</b>	2.8
<b>Among women with unintended pregnancy, % using conception at conception</b>			
	1998-2003	<b>46.7</b>	1.9
	1998	<b>39.7</b>	5.7
	1999	<b>44.5</b>	4.4
	2000	<b>53.4</b>	4.2
	2001	<b>49.9</b>	4.4
	2002	<b>45.2</b>	4.1
	2003	<b>47.9</b>	4.6
<b>Among women not trying to get pregnant, % using conception at conception</b>			
	2000-2003	<b>44.5</b>	1.8
	2000	<b>47.3</b>	3.5
	2001	<b>44.4</b>	3.8
	2002	<b>42.6</b>	3.5
	2003	<b>43.7</b>	3.9
<b>Among women with unintended pregnancy, % not using conception at conception</b>			
	1998-2003	<b>53.3</b>	1.9
	1998	<b>60.4</b>	5.7
	1999	<b>55.5</b>	4.4
	2000	<b>46.7</b>	4.2
	2001	<b>50.1</b>	4.4
	2002	<b>54.8</b>	4.1
	2003	<b>52.1</b>	4.6
<b>Among women not trying to get pregnant, % not using preconception conception</b>			
	2000-2003	<b>55.5</b>	1.8
	2000	<b>52.7</b>	3.5
	2001	<b>55.6</b>	3.8
	2002	<b>57.4</b>	3.5
	2003	<b>56.3</b>	3.9
<b>% who were using postpartum contraception</b>			
	1998-2003	<b>84.8</b>	0.9
	1998	<b>78.7</b>	2.9
	1999	<b>82.5</b>	2.1
	2000	<b>86.9</b>	1.7
	2001	<b>85.2</b>	1.9
	2002	<b>88.6</b>	1.7
	2003	<b>86.7</b>	2.0

	Year of birth	Percent	±
<b>% who drank any alcohol during the 3 months before pregnancy</b>			
	1998-2003	<b>46.7</b>	1.2
	1998	<b>45.6</b>	3.6
	1999	<b>45.4</b>	2.8
	2000	<b>46.9</b>	2.6
	2001	<b>48.5</b>	2.7
	2002	<b>46.5</b>	2.6
	2003	<b>47.3</b>	2.8
<b>% who drank any alcohol during the last 3 months of pregnancy</b>			
	1998-2003	<b>4.6</b>	0.5
	1998	<b>4.6</b>	1.4
	1999	<b>4.3</b>	1.1
	2000	<b>5.2</b>	1.2
	2001	<b>4.6</b>	1.0
	2002	<b>4.7</b>	1.1
	2003	<b>4.1</b>	1.1
<b>% who drank alcohol frequently or binged during the 3 months before pregnancy</b>			
	1998-2003	<b>18.2</b>	0.9
	1998	<b>20.3</b>	2.9
	1999	<b>18.4</b>	2.2
	2000	<b>17.8</b>	2.0
	2001	<b>19.0</b>	2.2
	2002	<b>17.4</b>	2.0
	2003	<b>16.7</b>	2.1
<b>% who drank alcohol frequently or binged during the last 3 months of pregnancy</b>			
	1998-2003	<b>1.0</b>	0.2
	1998	<b>1.2</b>	0.8
	1999	<b>0.6</b>	0.4
	2000	<b>1.1</b>	0.6
	2001	<b>0.7</b>	0.4
	2002	<b>1.0</b>	0.5
	2003	<b>1.2</b>	0.7

	Year of birth	Percent	±
<b>% who smoked any cigarettes during the 3 months before pregnancy</b>			
	1998-2003	<b>23.2</b>	1.0
	1998	<b>25.9</b>	3.2
	1999	<b>27.1</b>	2.5
	2000	<b>22.2</b>	2.2
	2001	<b>24.4</b>	2.4
	2002	<b>20.4</b>	2.1
	2003	<b>19.0</b>	2.2
<b>% who smoked any cigarettes during the last 3 months of pregnancy</b>			
	1998-2003	<b>10.3</b>	0.7
	1998	<b>12.6</b>	2.5
	1999	<b>11.6</b>	1.8
	2000	<b>9.7</b>	1.5
	2001	<b>10.7</b>	1.7
	2002	<b>8.5</b>	1.5
	2003	<b>8.5</b>	1.6
<b>% who currently smoke any cigarettes</b>			
	1998-2003	<b>16.5</b>	0.9
	1998	<b>19.5</b>	3.0
	1999	<b>20.1</b>	2.3
	2000	<b>16.4</b>	1.9
	2001	<b>16.2</b>	2.1
	2002	<b>13.7</b>	1.8
	2003	<b>12.8</b>	1.9
<b>% who said their infant is exposed to tobacco smoke</b>			
	1998-2003	<b>8.0</b>	0.7
	1998	<b>10.8</b>	2.4
	1999	<b>6.1</b>	1.4
	2000	<b>8.6</b>	1.5
	2001	<b>9.7</b>	1.6
	2002	<b>6.9</b>	1.3
	2003	<b>6.1</b>	1.4

	Year of birth	Percent	±
<b>% whose partner physically abused them during the 12 months before pregnancy</b>			
	1998-2003	<b>7.6</b>	0.6
	1998	<b>8.4</b>	2.0
	1999	<b>7.7</b>	1.5
	2000	<b>8.2</b>	1.5
	2001	<b>6.0</b>	1.2
	2002	<b>7.6</b>	1.4
	2003	<b>7.6</b>	1.5
<b>% whose partner physically abused them during pregnancy</b>			
	1998-2003	<b>6.2</b>	0.6
	1998	<b>7.0</b>	1.8
	1999	<b>6.3</b>	1.4
	2000	<b>6.6</b>	1.3
	2001	<b>5.4</b>	1.1
	2002	<b>5.6</b>	1.2
	2003	<b>6.4</b>	1.4
<b>% who had a weight problem (at risk for overweight/overweight/obese) before pregnancy</b>	Birthyear		
	1998	<b>32.9</b>	3.3
	1999	<b>33.8</b>	2.6
	2000	<b>38.0</b>	2.5
	2001	<b>38.4</b>	2.6
	2002	<b>39.8</b>	2.6
	2003	<b>39.7</b>	2.8
<b>% who had pre-existing or gestational diabetes</b>			
	1998-2003	<b>7.7</b>	0.6
	1998	<b>7.0</b>	1.8
	1999	<b>7.4</b>	1.5
	2000	<b>7.3</b>	1.3
	2001	<b>7.6</b>	1.4
	2002	<b>8.6</b>	1.5
	2003	<b>8.3</b>	1.6

	Year of birth	Percent	±
<b>Prenatal care</b>			
<b>% who had late(after 1st trimester) or no prenatal care</b>			
	1998-2003	<b>30.9</b>	1.1
	1998	<b>35.1</b>	3.6
	1999	<b>32.3</b>	2.7
	2000	<b>30.4</b>	2.5
	2001	<b>29.6</b>	2.5
	2002	<b>28.9</b>	2.4
	2003	<b>29.3</b>	2.7
<b>% who had timely (within 1st trimester) prenatal care</b>			
	1998-2003	<b>69.1</b>	1.1
	1998	<b>64.9</b>	3.6
	1999	<b>67.7</b>	2.7
	2000	<b>69.6</b>	2.5
	2001	<b>70.4</b>	2.5
	2002	<b>71.1</b>	2.4
	2003	<b>70.7</b>	2.7
<b>% who had an adequate level of prenatal care (Kotelchuck index)</b>			
	1998-2003	<b>33.4</b>	1.1
	1998	<b>33.5</b>	3.5
	1999	<b>34.6</b>	2.7
	2000	<b>33.0</b>	2.5
	2001	<b>30.7</b>	2.5
	2002	<b>33.4</b>	2.5
	2003	<b>35.4</b>	2.8
<b>Among women with late or no prenatal care, % who started as early as wanted</b>			
	1998-2003	<b>57.9</b>	2.3
	1998	<b>56.3</b>	6.7
	1999	<b>56.7</b>	5.2
	2000	<b>58.0</b>	5.1
	2001	<b>55.4</b>	5.0
	2002	<b>60.9</b>	4.9
	2003	<b>60.6</b>	5.6

	Year of birth	Percent	±
<i>Among women who had late or no prenatal care and did not start as early as desired, % with these reasons for late entry:</i>			
<b>Did not know she was pregnant</b>			
	1998-2003	<b>34.8</b>	3.4
	1998-1999	<b>32.3</b>	6.3
	2000-2003	<b>36.2</b>	4.0
<b>Lacked money or insurance</b>			
	1998-2003	<b>32.0</b>	3.4
	1998-1999	<b>34.9</b>	6.6
	2000-2003	<b>30.3</b>	3.8
<b>Was unable to get an appointment</b>			
	1998-2003	<b>25.3</b>	3.1
	1998-1999	<b>24.9</b>	5.9
	2000-2003	<b>25.5</b>	3.6
<b>Had too many other things going on</b>			
	1998-2003	<b>17.5</b>	2.5
	1998-1999	<b>16.0</b>	4.3
	2000-2003	<b>18.4</b>	3.0
<b>Lacked Medicaid card</b>			
	2000-2003	<b>18.4</b>	3.1
<b>Lacked transportation to clinic/office</b>			
	1998-2003	<b>11.6</b>	2.3
	1998-1999	<b>9.8</b>	4.2
	2000-2003	<b>12.6</b>	2.6
<b>Had other reasons</b>			
	1998-2003	<b>13.9</b>	2.6
	1998-1999	<b>15.4</b>	5.1
	2000-2003	<b>13.0</b>	2.8
<b>Was delayed by doctor or health plan</b>			
	2000-2003	<b>6.3</b>	2.0
<b>Could not get child care</b>			
	1998-2003	<b>4.2</b>	1.3
	1998-1999	<del><b>2.8</b></del>	<del>4.9</del>
	2000-2003	<b>5.0</b>	1.7

	Year of birth	Percent	±
<b>% who recalled discussion of these topics with a prenatal healthcare worker</b>			
<b>Breastfeeding</b>			
	1998-2003	<b>87.4</b>	0.8
	1998	<b>88.5</b>	2.3
	1999	<b>88.3</b>	1.8
	2000	<b>85.8</b>	1.8
	2001	<b>87.5</b>	1.8
	2002	<b>87.1</b>	1.8
	2003	<b>87.1</b>	1.9
<b>Medicines that are safe during pregnancy</b>			
	1998-2003	<b>87.4</b>	0.8
	1998	<b>88.4</b>	2.3
	1999	<b>88.6</b>	1.8
	2000	<b>86.2</b>	1.8
	2001	<b>86.4</b>	1.9
	2002	<b>86.6</b>	1.8
	2003	<b>88.2</b>	1.9
<b>Birth control methods to use after pregnancy</b>			
	1998-2003	<b>85.1</b>	0.8
	1998	<b>83.9</b>	2.7
	1999	<b>87.3</b>	1.9
	2000	<b>85.5</b>	1.8
	2001	<b>82.8</b>	2.1
	2002	<b>86.3</b>	1.8
	2003	<b>84.8</b>	2.0
<b>Doing tests to screen for birth defects in the family</b>			
	2000-2003	<b>83.7</b>	1.0
	2000	<b>81.4</b>	2.1
	2001	<b>82.2</b>	2.1
	2002	<b>85.1</b>	1.9
	2003	<b>86.2</b>	2.0
<b>What to do for early labor</b>			
	1998-2003	<b>84.3</b>	0.8
	1998	<b>85.4</b>	2.5
	1999	<b>87.7</b>	1.8
	2000	<b>82.9</b>	1.9
	2001	<b>80.9</b>	2.1
	2002	<b>83.3</b>	1.9
	2003	<b>85.8</b>	2.0

	Year of birth	Percent	±
<b>% who recalled discussion of these topics with a prenatal healthcare worker</b>			
<b>Getting a blood test for HIV</b>			
	1998-2003	<b>80.1</b>	0.9
	1998	<b>77.2</b>	3.0
	1999	<b>78.5</b>	2.3
	2000	<b>78.5</b>	2.2
	2001	<b>80.8</b>	2.1
	2002	<b>82.8</b>	2.0
	2003	<b>82.5</b>	2.2
<b>How maternal alcohol use during pregnancy could affect the baby</b>			
	1998-2003	<b>77.0</b>	1.0
	1998	<b>82.0</b>	2.7
	1999	<b>84.6</b>	2.0
	2000	<b>74.3</b>	2.3
	2001	<b>72.8</b>	2.4
	2002	<b>74.9</b>	2.3
	2003	<b>73.8</b>	2.6
<i>Among women who drank alcohol during late pregnancy, % who recalled discussion about how maternal alcohol use during pregnancy could affect the baby</i>			
	1998-2003	<b>75.4</b>	4.7
	1998-1999	<b>79.6</b>	4.5
	2000-2003	<b>73.3</b>	5.3
<b>How maternal smoking during pregnancy could affect the baby</b>			
	1998-2003	<b>75.7</b>	1.0
	1998	<b>81.8</b>	2.8
	1999	<b>83.0</b>	2.1
	2000	<b>71.7</b>	2.3
	2001	<b>71.8</b>	2.5
	2002	<b>74.3</b>	2.3
	2003	<b>71.9</b>	2.6
<i>Among women who smoked during late pregnancy, % who recalled discussion about how maternal smoking could affect the baby</i>			
	1998-2003	<b>87.1</b>	2.5
	1998-1999	<b>93.1</b>	3.5
	2000-2003	<b>83.3</b>	3.4

	Year of birth	Percent	±
<b>% who recalled discussion of these topics with a prenatal healthcare worker</b>			
<b>How illegal drugs could affect the baby</b>			
	1998-2003	<b>70.5</b>	1.1
	1998	<b>75.7</b>	3.1
	1999	<b>77.9</b>	2.3
	2000	<b>66.2</b>	2.5
	2001	<b>66.5</b>	2.6
	2002	<b>69.0</b>	2.5
	2003	<b>67.9</b>	2.7
<b>Using a seat belt during pregnancy</b>			
	1998-2003	<b>59.1</b>	1.2
	1998	<b>61.8</b>	3.5
	1999	<b>63.3</b>	2.7
	2000	<b>55.7</b>	2.6
	2001	<b>55.4</b>	2.7
	2002	<b>57.1</b>	2.6
	2003	<b>61.3</b>	2.8
<b>Physical abuse to women by their husband or partner</b>			
	1998-2003	<b>45.2</b>	1.2
	1998	<b>37.4</b>	3.5
	1999	<b>43.2</b>	2.8
	2000	<b>42.5</b>	2.6
	2001	<b>44.6</b>	2.7
	2002	<b>51.3</b>	2.7
	2003	<b>52.1</b>	2.9
<i>Among women whose partner abused them during pregnancy, % who recalled discussion about <b>partner abuse</b></i>			
	1998-2003	<b>49.5</b>	4.8
	1998-1999	<b>47.2</b>	<del>9.0</del>
	2000-2003	<b>50.8</b>	5.5
<b>How to care for her teeth and gums</b>			
	1998-2003	<b>20.3</b>	0.9
	1998	<b>13.0</b>	2.4
	1999	<b>16.8</b>	2.1
	2000	<b>22.2</b>	2.1
	2001	<b>20.3</b>	2.1
	2002	<b>24.6</b>	2.3
	2003	<b>24.6</b>	2.4

	Year of birth	Percent	±
<b>Payer of healthcare and services</b>			
<b>Preconception healthcare payer</b>			
<b>% of women with...</b>			
Medicaid			
	1998-2003	<b>15.8</b>	0.9
	1998	<b>11.9</b>	2.4
	1999	<b>13.1</b>	1.9
	2000	<b>14.7</b>	1.9
	2001	<b>16.4</b>	2.0
	2002	<b>18.3</b>	2.0
	2003	<b>20.6</b>	2.3
Insurance			
	1998-2003	<b>45.2</b>	1.2
	1998	<b>48.3</b>	3.5
	1999	<b>46.0</b>	2.8
	2000	<b>44.5</b>	2.5
	2001	<b>44.2</b>	2.6
	2002	<b>43.5</b>	2.5
	2003	<b>44.8</b>	2.8
<b>Prenatal healthcare payer: % of women with...</b>			
Medicaid			
	1998-2003	<b>48.8</b>	1.2
	1998	<b>44.4</b>	3.6
	1999	<b>49.9</b>	2.8
	2000	<b>48.1</b>	2.6
	2001	<b>49.4</b>	2.7
	2002	<b>48.3</b>	2.6
	2003	<b>52.9</b>	2.8
Insurance			
	1998-2003	<b>41.8</b>	1.1
	1998	<b>45.3</b>	3.5
	1999	<b>42.9</b>	2.8
	2000	<b>42.3</b>	2.5
	2001	<b>41.5</b>	2.6
	2002	<b>40.3</b>	2.5
	2003	<b>38.4</b>	2.7
Indian Health Service			
	1998-2003	<b>7.1</b>	0.5
	1998	<b>8.3</b>	0.7
	1999	<b>8.1</b>	1.1
	2000	<b>7.4</b>	1.4
	2001	<b>6.5</b>	1.0
	2002	<b>6.9</b>	1.2
	2003	<b>5.8</b>	1.2
Medicaid or insurance or IHS			
	1998-2003	<b>88.3</b>	0.8
	1998	<b>88.0</b>	2.5
	1999	<b>90.8</b>	1.7
	2000	<b>89.1</b>	1.7
	2001	<b>88.2</b>	1.8
	2002	<b>86.5</b>	1.8
	2003	<b>87.4</b>	1.9

	Year of birth	Percent	±
<b>Payer of delivery</b>			
<b>% of women with...</b>			
Medicaid			
	1998-2003	<b>53.5</b>	1.2
	1998	<b>48.0</b>	3.6
	1999	<b>52.2</b>	2.8
	2000	<b>54.5</b>	2.6
	2001	<b>53.9</b>	2.6
	2002	<b>54.3</b>	2.6
	2003	<b>58.2</b>	2.8
Insurance			
	1998-2003	<b>39.8</b>	1.1
	1998	<b>43.4</b>	3.5
	1999	<b>39.9</b>	2.7
	2000	<b>41.0</b>	2.5
	2001	<b>40.5</b>	2.6
	2002	<b>37.8</b>	2.5
	2003	<b>36.5</b>	2.7
Indian Health Service			
	1998-2003	<b>5.2</b>	0.4
	1998	<b>6.8</b>	0.7
	1999	<b>6.1</b>	1.0
	2000	<b>4.9</b>	1.1
	2001	<b>4.6</b>	0.9
	2002	<b>5.0</b>	1.0
	2003	<b>3.9</b>	1.0
Medicaid or insurance or IHS			
	1998-2003	<b>92.1</b>	0.7
	1998	<b>91.3</b>	2.1
	1999	<b>92.5</b>	1.5
	2000	<b>93.5</b>	1.3
	2001	<b>92.2</b>	1.5
	2002	<b>90.9</b>	1.5
	2003	<b>92.1</b>	1.5

	Year of birth	Percent	±
<b>Services during and after pregnancy</b>			
<b>% of women who had</b>			
WIC during pregnancy (core question)			
	1998-2003	<b>55.8</b>	1.2
	1998	<b>58.4</b>	3.5
	1999	<b>56.1</b>	2.8
	2000	<b>54.7</b>	2.6
	2001	<b>53.0</b>	2.6
	2002	<b>55.3</b>	2.6
	2003	<b>57.5</b>	2.8
WIC during pregnancy (core/PGSV)			
	1998-2003	<b>58.0</b>	1.1
	1998	<b>60.5</b>	3.4
	1999	<b>58.1</b>	2.7
	2000	<b>57.6</b>	2.5
	2001	<b>55.4</b>	2.6
	2002	<b>57.1</b>	2.6
	2003	<b>59.1</b>	2.8
Prenatal WIC classes or groups			
	1998-2003	<b>41.8</b>	1.1
	1998	<b>42.1</b>	3.5
	1999	<b>44.0</b>	2.8
	2000	<b>42.7</b>	2.6
	2001	<b>39.5</b>	2.5
	2002	<b>41.4</b>	2.5
	2003	<b>41.1</b>	2.8
Postpartum WIC classes or groups			
	1998-2003	<b>38.8</b>	1.1
	1998	<b>35.5</b>	3.4
	1999	<b>36.2</b>	2.7
	2000	<b>42.3</b>	2.6
	2001	<b>38.0</b>	2.6
	2002	<b>40.2</b>	2.5
	2003	<b>40.4</b>	2.8

	Year of birth	Percent	±
<b>Services during and after pregnancy</b>			
<b>% of women who had</b>			
Prenatal home visiting services			
	1998-2003	<b>6.2</b>	0.6
	1998	<b>7.6</b>	1.9
	1999	<b>10.5</b>	1.7
	2000	<b>4.6</b>	1.0
	2001	<b>4.9</b>	1.1
	2002	<b>5.3</b>	1.2
	2003	<b>4.4</b>	1.0
Postpartum home visiting services			
	1998-2003	<b>10.6</b>	0.7
	1998	<b>12.5</b>	2.2
	1999	<b>17.6</b>	2.1
	2000	<b>9.4</b>	1.5
	2001	<b>8.2</b>	1.4
	2002	<b>8.9</b>	1.5
	2003	<b>7.1</b>	1.4
<i>Among women with Medicaid for prenatal care, % with prenatal Families FIRST</i>			
	1998-2003	<b>12.2</b>	1.1
	2000	<b>12.5</b>	2.5
	2001	<b>14.1</b>	2.3
	2002	<b>12.6</b>	2.3
	2003	<b>9.7</b>	2.0
<i>Among women with Medicaid for delivery, % with postpartum Families FIRST</i>			
	1998-2003	<b>8.7</b>	0.9
	2000	<b>8.8</b>	1.9
	2001	<b>9.6</b>	1.9
	2002	<b>7.8</b>	1.7
	2003	<b>8.7</b>	1.8

	Year of birth	Percent	±
<b>Services during and after pregnancy</b>			
<b>% of women who had</b>			
Prenatal breastfeeding classes or groups			
	1998-2003	<b>17.0</b>	0.9
	1998	<b>18.2</b>	2.7
	1999	<b>21.3</b>	2.2
	2000	<b>16.5</b>	1.9
	2001	<b>15.3</b>	1.8
	2002	<b>15.4</b>	1.8
	2003	<b>15.4</b>	2.0
Postpartum breastfeeding classes or groups			
	1998-2003	<b>10.3</b>	0.7
	1998	<b>8.9</b>	1.9
	1999	<b>11.3</b>	1.7
	2000	<b>11.6</b>	1.6
	2001	<b>10.5</b>	1.6
	2002	<b>10.5</b>	1.6
	2003	<b>8.8</b>	1.6
Prenatal parenting classes or groups			
	1998-2003	<b>18.4</b>	0.9
	1998	<b>19.6</b>	2.8
	1999	<b>20.4</b>	2.2
	2000	<b>17.2</b>	1.9
	2001	<b>17.8</b>	2.0
	2002	<b>17.6</b>	2.0
	2003	<b>17.5</b>	2.2
Postpartum parenting classes or groups			
	1998-2003	<b>4.9</b>	0.5
	1998	<b>3.6</b>	1.3
	1999	<b>6.6</b>	1.4
	2000	<b>4.9</b>	1.1
	2001	<b>4.3</b>	1.0
	2002	<b>5.8</b>	1.2
	2003	<b>4.2</b>	1.1

	Year of birth	Percent	±
<b>Services during and after pregnancy: special groups</b>			
<i>Among women who might benefit from counseling, % with prenatal counseling</i>			
	1998-2003	<b>6.5</b>	0.6
	1998-1999	<b>8.2</b>	1.2
	2000-2003	<b>5.6</b>	0.6
<i>Among women who might benefit from counseling, % with postpartum counseling</i>			
	1998-2003	<b>6.8</b>	0.9
	1998-1999	<b>7.7</b>	1.7
	2000-2003	<b>6.4</b>	1.0
<i>Among women physically abused by their partner 12 mo. before or during pregnancy, % who used a prenatal program for protection from family violence</i>			
	1998-2003	<b>5.9</b>	2.0
	1998-1999	<del>6.3</del>	<del>3.8</del>
	2000-2003	<b>5.7</b>	2.3
<i>Among women physically abused by their partner during pregnancy, % who used a postpartum program for protection from family violence</i>			
	1998-2003	<b>6.8</b>	2.6
	1998-1999	<b>8.9</b>	5.0
	2000-2003	<b>5.6</b>	2.8
<i>Among women who drank frequently or binged during the 3 months before pregnancy, % who used a prenatal program for drug or alcohol cessation</i>			
	1998-2003	<b>2.0</b>	0.8
	1998-1999	<b>3.3</b>	1.9
	2000-2003	<del>4.3</del>	<del>0.8</del>
<i>Among women who smoked 3 months before or during pregnancy, % who joined prenatal smoking cessation classes or groups</i>			
	1998-2003	<b>2.3</b>	0.8
	1998-1999	<b>3.6</b>	1.7
	2000-2003	<b>1.4</b>	0.7
<i>Among women who smoked during last 3 months of pregnancy or currently, % who joined postpartum smoking cessation classes or groups</i>			
	1998-2003	<b>1.0</b>	0.5
	1998-1999	<del>4.2</del>	<del>0.9</del>
	2000-2003	<del>0.9</del>	<del>0.7</del>
<i>Among pregnant teens, % with prenatal services for teens</i>			
	1998-2003	<b>18.4</b>	2.3
	1998-1999	<b>21.0</b>	4.5
	2000-2003	<b>17.0</b>	2.5
<i>Among parenting teens, % with postpartum services for teens</i>			
	1998-2003	<b>9.0</b>	1.6
	1998-1999	<b>10.2</b>	3.3
	2000-2003	<b>8.4</b>	1.8

	Year of birth	Percent	±
<b>Oral health</b>			
% who recalled discussion with a prenatal healthcare worker about how to care for her teeth and gums			
	1998-2003	<b>20.3</b>	0.9
	1998	<b>13.0</b>	2.4
	1999	<b>16.8</b>	2.1
	2000	<b>22.2</b>	2.1
	2001	<b>20.3</b>	2.1
	2002	<b>24.6</b>	2.3
	2003	<b>24.6</b>	2.4
% who had a dental problem			
	1998-2003	<b>13.0</b>	0.8
	1998	<b>11.9</b>	2.3
	1999	<b>14.2</b>	2.0
	2000	<b>12.3</b>	1.7
	2001	<b>13.2</b>	1.8
	2002	<b>12.9</b>	1.8
	2003	<b>13.5</b>	2.0
% who had dental care			
	1998-2003	<b>28.3</b>	1.0
	1998	<b>21.5</b>	2.9
	1999	<b>22.7</b>	2.3
	2000	<b>28.7</b>	2.3
	2001	<b>30.2</b>	2.5
	2002	<b>33.2</b>	2.5
	2003	<b>33.7</b>	2.7
<i>Among women with a dental problem , % who received dental care</i>			
	1998-2003	<b>44.9</b>	3.2
	1998	<b>42.0</b>	<del>40.4</del>
	1999	<b>36.4</b>	7.2
	2000	<b>47.6</b>	7.3
	2001	<b>43.1</b>	7.3
	2002	<b>55.9</b>	7.2
	2003	<b>45.0</b>	<del>7.7</del>

	Year of birth	Percent	±
<b>Infant care, nutrition</b>			
Percent of women who initiated breastfeeding			
	1998-2003	<b>79.7</b>	1.0
	1998	<b>75.8</b>	3.2
	1999	<b>78.1</b>	2.4
	2000	<b>79.9</b>	2.1
	2001	<b>80.1</b>	2.2
	2002	<b>82.0</b>	2.1
	2003	<b>82.4</b>	2.2
Percent of all women who were breastfeeding when infant was at least 9 weeks old			
	1998-2003	<b>53.5</b>	1.2
	1998	<b>49.0</b>	3.6
	1999	<b>47.1</b>	2.8
	2000	<b>55.4</b>	2.6
	2001	<b>55.1</b>	2.7
	2002	<b>57.7</b>	2.6
	2003	<b>56.9</b>	2.9
Among women who initiated breastfeeding, % who continued breastfeeding until infant was at least 9 weeks old			
	1998-2003	<b>67.2</b>	1.2
	1998	<b>64.6</b>	3.9
	1999	<b>60.3</b>	3.1
	2000	<b>69.4</b>	2.7
	2001	<b>68.8</b>	2.8
	2002	<b>70.4</b>	2.7
	2003	<b>69.0</b>	3.0
Percent of women who place their infant to sleep on his/her back			
	1998-2003	<b>59.0</b>	1.2
	1998	<b>45.3</b>	3.6
	1999	<b>53.6</b>	2.9
	2000	<b>60.2</b>	2.6
	2001	<b>66.3</b>	2.5
	2002	<b>63.9</b>	2.5
	2003	<b>64.8</b>	2.7
Percent of women whose infant had an adequate number of well-baby visits			
	1998-2003	<b>79.8</b>	1.0
	1998	<b>82.3</b>	2.8
	1999	<b>82.0</b>	2.2
	2000	<b>81.1</b>	2.1
	2001	<b>78.3</b>	2.3
	2002	<b>76.5</b>	2.3
	2003	<b>78.4</b>	2.5
Percent of women whose family had enough food			
	2000-2003	<b>84.9</b>	1.0
	2000	<b>84.1</b>	1.9
	2001	<b>86.0</b>	1.9
	2002	<b>86.0</b>	1.8
	2003	<b>83.3</b>	2.2